








Nasbíráno 1.4.2002 pro přepsání do dvouznakového zápisu




$$B = 1 + 1 \neq 1 + 0 + 0$$



$$\frac{q}{e} = m_I + \frac{S+B}{2}$$

$m_I$  = projection of isospin  
 $S$  = strangeness  
 $B$  = baryon number



$$I = 1 \neq 1/2 + 0 \text{ Isospin}$$




$U$  = "up" quark  $+\frac{2}{3}e$   
 $D$  = "down" quark  $-\frac{1}{3}e$

$m_p = 1836.15 m_e$   
 Mass =  $1.6726 \times 10^{-27} \text{ kg}$   
 $= 938.27231 \text{ MeV}/c^2$   
 $= 1.00727647 \text{ u}$

composed of two down [quarks](#) and one up quark.

A free neutron will decay with a [half-life](#) of about 10.3 minutes but it is stable if combined into a nucleus. The decay of the neutron involves the [weak interaction](#) as indicated in the [Feynman diagram](#) to the right. This fact is important in models of the [early universe](#). The neutron is about 0.2% more massive than a proton, which translates to an energy difference of 1.29 MeV.



$U$  = "up" quark  $+\frac{2}{3}e$   
 $D$  = "down" quark  $-\frac{1}{3}e$

$m_p = 1838.68 m_e$   
 Mass =  $1.6749 \times 10^{-27} \text{ kg}$   
 $= 939.5656 \text{ MeV}/c^2$   
 $= 1.0086647 \text{ u}$

